

Exploring hydrocarbon degrading ability of bacteria isolated from engine oil polluted soil.

Chaithra. J. Rai¹, Bharathi Prakash², Girisha S K³

¹PG center for Microbiology, Mangalore University, Chikkalauvara, Kodagu.

² Department of Microbiology, University College, Mangalore University Hampankatta, Mangalore

³ College of Fisheries, KVAFSU, Mangalore

Article History

Received: 25/07/2023

Accepted: 15/08/2023

Article ID: RRBB/102

Corresponding Author:

E-Mail:

bharatiprakash21@gmail.com

Abstract:

This study was conducted to assess the capabilities of bacterial isolates obtained from oil-polluted sites with waste engine oil. Engine oil is made of crude petroleum, it mainly consists of hydrocarbons which includes aliphatic, aromatic hydrocarbons, alkyl compounds, resins etc. and trace amounts of Sulphur or nitrogen. To treat these toxic pollutants bioremediation is an environment friendly approach to reduce their intensity of toxicity.

Waste engine oil contaminated soil samples were collected from specific garage areas in Mangalore, and assessed for high hydrocarbon and oil degradation capacity. Isolates were processed for biochemical and molecular tests.

Emulsification index and gravimetric analysis was done to evaluate the isolates' efficacy in emulsifying and degrading potential of oil. Molecular tests confirmed the presence of three significant bacterial species – *Klebsiella quasipneumoniae*, *Stutzerimonas stutzeri*, and *Klebsiella pneumoniae* showing their potential for waste engine oil bioremediation.

In conclusion, this study uncovers the potential of bacterial isolates for effective waste engine oil biodegradation. It emphasizes the significance of employing these isolates to promote sustainable waste management practices, highlighting the need for further research to optimize bioremediation techniques and facilitate practical implementation in the field.

Keywords: Engine oil, Biodegradation, Emulsification index, Gravimetric analysis, bioremediation.

Introduction:

Waste engine oil, a significant byproduct of machinery and vehicle operation, contaminates surrounding environments like soil and water. (1) Treating used motor oil is a challenge because it doesn't mix with water, has a thick texture, and may contain harmful substances. It breaks down slowly and sticks to surfaces, resulting in pollution of water and soil. It imposes great risk to the environment and various forms of life. Bioremediation provides a cost-effective and efficient solution for cleaning up contaminated environments. However, careful microbial handling and continuous monitoring are necessary to address concerns in this approach. (2) By implementing proper waste disposal methods and preventing soil contamination, we can mitigate the risks it poses to ecosystems and human health.

Physical remediation, incineration and landfilling have minimal to no negative impact on marine biota, making them a sustainable choice. However, their effectiveness is influenced by various environmental and oceanic factors, including wave action, temperature, oil type, and the presence of natural dispersants. (3)

As a result, there is a growing need for sustainable and cost-effective approaches to tackle these environmental issues. (4) Bioremediation, using indigenous bacteria

Materials and Methods:

Sample collection: soil samples from the subsurface layer of soil from a depth of 5-30cm deep were collected from few engine oil contaminated sites.

and transferred catabolic genes, offers a targeted and effective solution for waste engine oil treatment. Isolating native petroleum-degrading strains and screening enzymes for over-expression through proteomic identification are crucial steps. This approach holds promise for addressing petroleum contamination. (5)

Factors such as oil composition, dispersion, and absorption, as well as temperature, oxygen, and nutrients, influence biodegradation rates. Bacteria and fungi are key degraders, with adaptation and genetic amplification enhancing degradation. Seeding can accelerate biodegradation under controlled conditions. (6)

This research explores the potential of bacterial isolates in efficiently and sustainably remediating waste engine oil.(7) Bacteria are isolated from oil-polluted sites, characterized, and tested for their oil degradation capabilities.(8) Molecular tests confirm the presence and identity of oil degrading bacterial species. The findings enhance our understanding of microbial diversity and functionality in oil-contaminated environments, paving the way for scalable bioremediation strategies. Effective waste engine oil treatment is crucial to mitigate environmental pollution and health risks associated with oil spills and contamination

1gm of the soil sample was added to 9ml of saline. It was vortexed and inoculated to Bushnell Hass (BH) broth supplemented with 2% engine oil as carbon source. After a brief growth period of 7 days the inoculums from this broth was spread plated on Bushnell Hass growth medium supplemented with engine oil as carbon source. (9,10, 11).

The plates were incubated at 37°C for 48 hrs. (12) The bacterial isolates obtained from the BH agar medium showed consistent growth as BH medium serves as a selective medium,

A. Biochemical characterization: Biochemical tests were performed to further identify and characterize the bacterial isolates. The tests are given in the table 1. Biochemical test results helped in classifying the isolates into relevant taxonomic groups. (13)

B. Oil degradation screening analysis. (14): Oil degrading capabilities of the The The

specifically designed to promote the growth of oil-degrading bacteria. (9)

Furthermore, after a thorough examination of the bacterial colonies on BH medium, distinct morphological characteristics such as variations in color, size, and shape were observed. Colonies representing unique characteristics were selected for further analysis and investigation. The pure cultures of the selected bacteria, individual colonies on BH media are shown in Figure.1

isolated bacteria were assessed using screening assays for their oil degrading ability. These assays involved the use of engine oil or specific hydrocarbon compounds as the sole carbon source. The degradation activity included changes in its Optical density, emulsification index and weight loss using analytical techniques such as gravimetric analysis

Figure1: Generation of pure bacterial cultures by sub culturing technique



Table: 1 Details of identified bacterial isolates and quantified results of gravimetric analysis.

Bacterial Isolate	Amount of oil degraded (gms)	Bacteria identified by 16s RNA Sequencing	Morphological features
CN1	0.25	<i>Klebsiella</i>	Gram-negative, non-motile, and rod-shaped bacteria, non-motile
CN8	0.32		

CN9	0.2	<i>pneumoniae</i>	
CN7	0.15	<i>Klebsiella quasipneumoniae</i>	Gram-negative, non-motile, and rod-shaped bacteria, non-motile
CN14	0.23		
CN16	0.36		
CN17	0.48		
CN18	0.48		
CN19	0.49		
CN13	0.31	<i>Stutzerimonas stutzeri</i>	Gram-negative motile, bacillus
CN15	0.46		

Note- Results of gravimetric analysis showed that CN15, CN17, CN18, and CN19 possess high capacity for oil degradation with degradation rates above 0.4gms, while CN1, CN7, CN14, CN8, CN9, CN13, and CN16 exhibited moderate to lower abilities for oil degradation.

Table 2. Biochemical tests of the bacterial isolates

Biochemical Tests	<i>Klebsiella pneumonia</i>	<i>Klebsiella quasipneumoniae</i>	<i>Stutzerimonas Stutzeri</i>
Carbohydrate fermentation test	+	+	+
Indole Test	-	-	-
Methyl red Test	-	+	+
Voges Proskauer (VP) test	+	+	+
Citrate Test	+	+	+
Urease Test	+	+	+
Catalase Test	+	+	+
Oxidase Test	-	-	+

C. Molecular identification: For a more precise identification of the isolated bacteria, 16S rRNA sequencing was carried out to determine the bacterial genus and species. (15) Identified isolates were preserved for future use. (16)

Results

Based on 16S rRNA sequencing, 3 bacterial species identified from oil-contaminated samples were *Klebsiella quasipneumoniae*, *Stutzerimonas stutzeri*, and *Klebsiella pneumoniae*. The sequences were submitted

to Gene bank. The accession numbers of gene bank submission in NCBI portal are

OP597535, OP597647, OP597648, OP597654, OP597675. These bacterial species provide valuable insights about the microbial composition of the oil-polluted sites.

The bacterial isolates CN15, CN17, CN18, and CN19 have demonstrated remarkable ability in breaking down more than 0.40 grams of oil, outperforming other bacterial isolates. The oil degrading capacity and biochemical test results of the isolates is given in the table 1 and 2 .



Conclusion:

In conclusion, oil degrading potential of these microorganisms, can be leveraged to standardize the environment friendly bioremediation protocols that will address oil pollution of water and soil due to waste engine oil.

Declaration of competing interest: There is no conflict of interest

Acknowledgements: The authors express their gratitude to the University College, Mangalore University, Mangalore, Karnataka, India, for supporting this research work.

References

1. Kvenvolden KA, Cooper CK. Natural seepage of crude oil into the marine environment. *Geo-Marine Letters*. 2003 Dec 1;23(3-4):140–6.
2. Vishwakarma GS, Bhattacharjee G, Gohil N, Singh V. Current status, challenges and future of bioremediation. *Bioremediation of Pollutants*. 2020;403–15.
3. Dhaka A, Chattopadhyay P. A review on physical remediation techniques for treatment of marine oil spills. *Journal of Environmental Management*. 2021 Jun; 288:112428.
4. Orellana R, Cumsille A, Piña-Gangas P, Rojas C, Arancibia A, Donghi S, et al. Economic Evaluation of Bioremediation of Hydrocarbon-Contaminated Urban Soils in Chile. *Sustainability* [Internet]. 2022 Jan 1 [cited 2022 Dec 3];14(19):11854. Available from: <https://www.mdpi.com/2071-1050/14/19/11854>
5. French KE, Zhou Z, Terry N. Horizontal “gene drives” harness indigenous bacteria for bioremediation. *Scientific Reports*. 2020 Sep 15;10(1).
6. Abdel-Shafy HI, Mansour MSM. Microbial Degradation of Hydrocarbons in the Environment: An Overview. *Microbial Action on Hydrocarbons*. 2018;353–86.
7. Mishra S, Jyot J, Kuhad RC, Lal B. Evaluation of Inoculum Addition To Stimulate In Situ Bioremediation of Oily-Sludge-Contaminated Soil. *Applied and Environmental Microbiology* [Internet]. 2001 Apr 1;67(4):1675–81. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC92784/>
8. Kumar V, Singh S, Manhas A, Singh J, Singla S, Kaur P, et al. Bioremediation of Petroleum hydrocarbon by using *Pseudomonas* species isolated from Petroleum contaminated soil. *Oriental Journal of Chemistry*. 2014 Dec 31;30(4):1771–6. <http://dx.doi.org/10.13005/ojc/300436>
9. Bushnell LD, Haas HF. The Utilization of Certain Hydrocarbons by Microorganisms 1. *Journal of Bacteriology*. 1941;41(5):653–73.
10. Gupte A, Sonawadekar sheetal. Study of Oil Degrading Bacteria Isolated From Oil Contaminated Sites. 2005.

11. Leahy JG, Colwell RR. Microbial degradation of hydrocarbons in the environment. *Microbiological Reviews*. 1990 Sep;54(3):305-315. <https://doi.org/10.1128/mr.54.3.305-315.1990>
 12. Dasgupta D, Ghosh R, Sengupta TK. Biofilm-Mediated Enhanced Crude
 13. Chauhan A, Jindal Tanu. *Biochemical and Molecular Methods for Bacterial Identification*. 2020.
 14. Kalami Reyhaneh, Pourbabaee Ahmad Ali, Alkhani Hussain Ali. Investigating of emulsification index and the growth capability of isolated bacteria from oil-contaminated soils in mineral medium containing diesel oil.
 15. Franco-Duarte R, Černáková L, Kadam S, S. Kaushik K, Salehi B, Bevilacqua A, et al. Advances in Chemical and Biological Methods to Identify Microorganisms – From past to Present. *Microorganisms*. 2019 May 13;7(5):130. <https://doi.org/10.3390/microorganisms7050130>
 16. Prakash O, Nimonkar Y, Shouche YS. Practice and prospects of microbial preservation. *FEMS Microbiology Letters* [Internet]. 2012 Nov 22;339(1):1-9. Available from: <https://academic.oup.com/femsle/article/339/1/1/496578> <https://doi.org/10.1111/1574-6968.12034>
- Oil Degradation by Newly Isolated Pseudomonas Species. *ISRN Biotechnology* [Internet]. 2013; 2013:1-13. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4393046/> <https://doi.org/10.5402/2013/250749>