

Survival strategies in plant parasitic nematodes

¹Babita Kumari, ²Sujata and ³Vinod Kumar

^{1&2}Ph.D. Scholar, Department of Nematology, CCS Haryana Agricultural University, Hisar

³Assistant Scientist, Department of Nematology, CCS Haryana Agricultural University, Hisar

Article History

Received: 02/11/2020

Revised: 15/11/2020

Accepted: 29/11/2020

<http://doi.org/10.5281/zenodo.4308456>



*Corresponding Author:

E-Mail: bk12081995@gmail.com

Introduction

'Survival strategies' are usually taken to mean the responses of organisms to the particular biological and physical challenges that may result in their death. Nematodes face a number of such challenges i.e. biological challenges and physical challenges. Biological challenges include: running out of food, predation, pathogens and competition. Parasitic species face the additional problems of the immune, and other, responses of their hosts and the problems of infecting new hosts. Physical challenges include: temperature extremes, desiccation, high pressure, acid or alkaline conditions, osmotic and ionic stress, anoxia and exposure to various types of radiation. The responses of nematodes to a physical challenge, however, also involve morphological, physiological, biochemical and molecular mechanisms. There may also be an interaction between the physical and biological challenges faced by nematodes.

There are four responses

1. **Migratory response:** The nematode might move away from the extreme conditions to a less stressful environment that is more favourable to its growth and reproduction.
2. **Capacity adaptation:** The nematode may adapt to the challenge and so be able to grow and reproduce under extreme conditions.
3. **Timing mechanism:** The life cycle of plant parasitic nematodes (PPNs) may include timing mechanisms that enable the parasite to avoid seasonally stressful conditions and to synchronize its life cycle with that of its host.
4. **Resistance adaptation:** The nematode may survive extremes in a dormant state until conditions

suitable for growth and reproduction return.

Adaptation strategies in plant-parasitic nematodes

Different PPNs also follow several survival strategies against environmental extremes and host response (Perry, 2011). Two types of dormancy, 'quiescence' and 'diapause' often lengthen survivability of unhatched larvae in soil living nematodes including root-knot nematode (Perry, 1989). Cryobiosis, thermobiosis, anoxybiosis, osmobiosis and desiccation are different survival mechanisms that are also observed in nematodes (Wright & Perry, 2006). First discovered PPNs, *Anguinatritici* have also special adaptation aptitude against desiccation (Needham, 1743). Commonly, life cycle of PPNs is divided into different distinct stages such as egg, four juvenile stages (J1, J2, J3 and J4) and adults and



different major PPN also exhibits different adaptation mechanism at their different life stages.

Eggs of root-knot nematode (RKN) are laid in mucoid protein mass that provides safety to eggs water loss and predators (Eisenback, 1985). Furthermore, extreme temperature makes this glycoprotein shriveled and hard that promotes mechanical pressure and hinders the hatching of J2 larva that is prone to drought environment (Wright & Perry, 2006). Additionally, gelatinous matrix defends the eggs from the attack of some soil microorganism as was observed in *M. javanica* (Orion & Kritzman, 2001). During dry conditions, a dried eggshell can also decrease water loss rate by changing its permeability (Ellenby, 1968). Additional research on eggshell permeability was done by Wharton (1980) and he observed that the lipid layer of the eggshell played a major role for lowering the water loss during desiccation. Not all PPN have survival capability. Mostly, the species with limited hosts such as potato cyst nematodes (*Globoderarostochiensis* and *G. pallida*) gained well adaption features for undesirable circumstances (Perry, 2002). The unhatched larvae of this nematode can survive many years in cyst without host (Turner & Rowe, 2006). Ellenby (1946) observed that the permeability of cyst wall exterior and eggshell of *G. rostochiensis* changed at desiccated conditions which helped cyst wall and eggshell to restrict their water losing rate compared to favourable conditions. This helps the unhatched susceptible J2 to survive in the egg and hatching factors (host root exudates) are obligatory for this nematode to hatch their J2 (Wright & Perry, 2006). J2 of soybean cyst nematode *Heteroderaglycines* hatched in water at optimum conditions but at the end of the plant growing season when environment is

becoming complex, most of the J2 remains in the cyst and hatching is solely dependent on hatching factors, irrespective to the origin of hatching factors; i.e. natural (Ishibashiet al., 1973) or artificial (Thompson & Tylka, 1997). Other cyst nematodes, such as *H. carotae* (Greco, 1981), *H. goettingiana* (Greco et al., 1986), *H. sacchari* (Ibrahim et al., 1993) contain the same characteristics. Even, RKN, *M. chitwoodi* and *M. triticoryzae* (Gaur et al., 2000; Wesemaelet al., 2006) also showed the same features for hatching of J2.

Under drying condition, species of *Anguina* and *Ditylenchus* showed coiling and clumping like features in their life cycle that helps to avoid water loss by reducing their surface (Crowe & Madin, 1975). *Ditylenchus myceliophagus* also lessen their water loss becoming coiled (Womersley, 1978). Furthermore, at the last stage of the crop growing season, *D. dipsaci*, stops their development at J4 stage due to shortage of food and make large aggregations and start coiling. This attribute known as 'eelworm wool' that helps them to remain alive for many years at desiccated condition by the sacrifice of peripheral J4s of this aggregation (Ellenby, 1969). Rice grain nematodes, *Aphelenchoides besseyi* also stops their multiplication at ripening stage of rice and adults become coiled and clumped which helps them surviving for 2-3 years with dry grains (Perry & Moens, 2011). On the other hand, J2 of 2 gall producing nematode *Anguinatritici* and *A. pacifica* become anhydrobiotic but remain uncoiled and survived many years (McClure et al., 2008). So, coiling is not obligatory for survival during desiccation. Nevertheless, *D. dipsaci* and *A. tritici*, can overcome desiccated environment and survive many years in above ground plant parts (Moens & Perry, 2009). Sheath retaining is also a survival mechanism against desiccated condition.

Young adults of *Rotylenchulus reniformis* remains confined with all 3 moulted cuticle of their larval stage which facilitates them of being dormant in adverse condition until situation becomes amiable for their survival (Gaur & Perry, 1991). Dauer phenomenon is a special characteristic of nematodes that helps them to survive at stress condition. Dauer is very common in *C. elegans* where dauer larvae restrict its biological activity and survive periods at desiccated situation (Kenyon, 1997). J2, J3, or J4 stages are performed as dauer in *A. tritici*, *C. elegans* and *D. dipsaci* respectively (Bird & Bird, 1991). Pine wood nematode *Bursaphelenchus xylophilus*, also contains a dauer stage that facilitates the transportation to pine trees with the help of insect vector *Monochamus* (Mota & Vieira, 2008).

Behavioural responses

Various types of aggregation, swarming and synchronized movements occur in nematodes. Some of these behaviours are associated with responses to environmental stress, including desiccation. Croll (1970) suggested that aggregation was associated with quiescence (reduced metabolism) and with survival responses to adverse environmental conditions, while swarming was involved in migration and dispersal. However, aggregation can occur in the absence of an environmental stress and swarming may be associated with such a stress. Aggregation as the clumping of nematodes either spontaneously in culture or in water or when they are drawn together by the evaporation of water to form a clump and Swarming as a coordinated movement by a mass of nematodes from one location to another. Swarming may be followed by aggregation. The best known example of aggregation is the accumulation of large

numbers of *D. dipsaci* on the surface of stored flower bulbs (Christie 1959). Croll (1970) mentions synchronized movements as another category and gives the movement of *Panagrellus redivivus* on to the walls of a culture vessel as an example.

Factors in survival of nematodes

"Hypobiosis" is the form of survival covering all the biological and metabolic states of an organism that are associated with inactive life and lowered metabolism.

"Dormancy" is the state of inactivity associated with a low but measurable metabolism, as contrasted with latent life or "cryptobiosis" which is associated with a nondetectable metabolism.

Dormancy (Hypometabolism)

It is a period in an organism's life cycle when growth, development, and (in animals) physical activity are temporarily stopped. This minimizes metabolic activity and therefore helps an organism to conserve energy. Dormancy tends to be closely associated with environmental conditions.

1. **Predictive dormancy** occurs when an organism enters a dormant phase before the onset of adverse conditions.
2. **Consequential dormancy** occurs when organisms enter a dormant phase after adverse conditions have arisen. It is considered to be a physiological state of dormancy with very specific initiating and inhibiting conditions.

Four types of dormancy in eggs of *Heterodera schachtii* is recognized (Zheng and Ferris 1991). Some eggs hatched quickly in water, some required host-root diffusate for rapid hatch, while others hatched slowly in water or in host-root diffusate. Stimulus

for hatching and ending of dormancy in various species include such factors as temperature (Van Gundy 1985) or the presence of host plant or root leachate (Huang and Pereira 1994; Sikora and Noel 1996). The quality of the latter depended on crop cultivar, phenology, and other factors (Sikora and Noel 1996). Interpretation of dormancy and diapause in nematode eggs is further complicated in that the induction of dormancy in cyst nematodes varies seasonally, and may be dependent on temperature or host phenology (Yen *et al.* 1995; Sikora and Noel 1996). Diapause and developmental dormancy seem to apply mostly to the egg stage and to juvenile stages within eggs, although instances of diapause in later juvenile stages or adults are known, mainly in a few animal-parasitic nematodes (Evans and Perry 1976).

Diapause is a mechanism used as a means to survive predictable, unfavorable environmental conditions, such as temperature extremes, drought, or reduced food availability. Diapause as a developmental arrest which is temporarily irreversible, so development will not resume, even under favourable conditions, until some intrinsic changes have been completed.

Resting stages in the life cycle

Infective stages: The life cycles of most parasitic nematodes include an infective stage in which development is arrested until infection occurs. The resting stage is usually the infective juvenile and this is often the stage of the life cycle that is most resistant to environmental stress.

Dauer juvenile formation: *Caenorhabditiselegans* produces a special survival stage called a dauer juvenile (dauer larva) in response to deteriorating environmental conditions. This is an

alternative developmental pathway that leads to the formation of dauer juveniles, rather than to normal J3 and reproductive development. E.g. infective juvenile (J2) of some anguinid plant-parasitic nematodes have also been referred to as dauer juveniles (Womersley *et al.*, 1998).

Diapause: It is defined as a period of developmental arrest that is either programmed into the life cycle (obligatory diapause) or is triggered by environmental conditions (facultative diapause). Diapause and other delays in developmental stages that are common in insects (Chapman 1971; Romoser & Stoffolano 1998) occur in some nematodes also (Evans & Perry 1976; Wharton 1986). Although diapause is not necessarily due to adverse environmental conditions nor ended by presence of favorable conditions (Chapman 1971; Evans & Perry 1976; Wharton 1986), it is nonetheless a critical survival mechanism during cold seasons and in the absence of a host. The increase of egg hatching in *Meloidogynenaasi* by chilling is a well-known example of diapause in a plant-parasitic nematodes (Van Gundy 1985). In some species of root-knot (*Meloidogyne* spp.) and cyst (*Heterodera* spp., *Globodera* spp.) nematodes, a portion of the eggs hatch rapidly while others hatch gradually over time (DeGuiran 1979; Zheng & Ferris 1991; Huang and Pereira 1994).

Egg diapause: Hatching of the eggs of the plant-parasitic nematodes *Heteroderaavenae* and *Meloidogynenaasi* is greater after chilling. The response to chilling varies among isolates of these nematodes and may allow overwintering or the survival of a dry season.

Quiescence: "Quiescence" most accurately describes dormancy in nematodes. The arrested development or movement of the

nematode during quiescence or cryptobiosis is usually caused by some unfavorable environmental conditions. It may be associated with a particular stage of development but is not necessarily associated with some type of morphogenesis. Quiescence is contrasted with diapause i.e., development will not resume unless or until certain physiological processes have been completed, even though the environment is favorable.

Cryptobiosis: Cryptobiosis is an ametabolic state of life entered by an organism in response to adverse environmental conditions such as desiccation, freezing, and oxygen deficiency. In the cryptobiotic state, all measurable metabolic processes stop, preventing reproduction, development, and repair. When environmental conditions return to being hospitable, the organism will return to its metabolic state of life as it was prior to the cryptobiosis. E.g. nematodes stored under dry conditions survive for long periods of time. *Tylenchus polyhynchus* was revived after 39 years, *Anguinatritici*, *Ditylenchus dipsaci* and *Tylenchus balsamophilus* after 20 years storage in plant tissue (Fielding 1951).

Chemobiosis is the cryptobiotic response to high levels of environmental toxins. It has been observed in tardigrades.

Cryobiosis is a form of cryptobiosis that takes place in reaction to decreased temperature. Cryobiosis initiates when the water surrounding the organism's cells has been frozen, stopping molecule mobility and allowing the organism to endure the freezing temperatures until more hospitable conditions return. The nematodes which are freezing-susceptible are those which escape freezing injury by supercooling at temperatures just below 0° C, but which freeze and die at some temperature.

The freezing-resistant nematodes are not injured by freezing. The group of nematodes considered susceptible to chilling injury may be further segregated into sensitive and tolerant.

Osmobiosis is the least studied of all types of cryptobiosis. Osmobiosis occurs in response to increased solute concentration in the solution the organism lives in. E.g. most nematodes are capable of tolerating and surviving at osmotic pressures up to 15-20 atm there is considerable variability in their tolerance to single ions. When the nematodes *T. Christiei*, *T. claytoni*, *H. erythrinae* (adults), *M. Javanica* and *T. semipenetrans* (larvae) were bathed in solutions of copper ions, they were still active, respectively, in the following solutions: 0 to 0.25, 0 to 4, 4 to 8, 16 to 32, >32 ppm for 24 hr.

Anhydrobiosis: Some nematodes can withstand moderate desiccation better than others, probably because the water distribution in the cells and the nature of the protein constituents are such that withdrawal of water does not cause the disruption of their intimate structure. *Ditylenchus dipsaci* larvae and *Trichostrongylus colubriformis* larvae have survived desiccation at 50 per cent relative humidity and 24° C for at least 32 days.

Survival responses to desiccation: For most nematodes, activity ceases once the surrounding water film evaporates. The loss of activity is due to the adherence of nematodes to the substrate in the absence of water or to the loss of turgor pressure, without which movement cannot occur. Nematodes may survive partial desiccation with a reduced rate of metabolism (desiccation-induced quiescence) or, after complete desiccation, metabolism may cease altogether (anhydrobiosis). A slow rate of

water loss is important for anhydrobiotic survival. *Anguinatritici*, *D. dipsaci* and *Tylenchus* have been showed potentiality of being enliven from desiccation after 20 year (Perry & Moens, 2011), 28 years (Fielding, 1951) and 39 years (Steiner & Albin, 1946) respectively. Cuticle lipid content of desiccated J4 of *D. dipsaci* and J2 of *A. tritici* have been increased compared to hydrated juveniles of those nematodes (Bird & Buttrose, 1974; Preston & Bird, 1987).

Anhydrobiotic nematodes can be divided into two groups: depending upon the rates of water loss from their environment that they will survive. Slow-dehydration strategists depend upon their surroundings, such as soil or moss, drying slowly; fast-dehydration strategists can survive rapid loss of water from their surroundings but themselves have mechanisms for ensuring the slow loss of water from their bodies. Adaptations for reducing the rate of water loss include both behavioural and physiological mechanisms (Table 1).

Table 1. Examples of nematode survival after lengthy time in anhydrobiosis

| Nematode | Normal active habits | Anhydrobiosis conditions | Time in anhydrobiosis | Reference |
|---------------------------------|-------------------------------|--------------------------|-----------------------|-----------------------------|
| <i>Anguinaagrostis</i> | Foliar plant parasite | Dried plant material | 4 yr | Fielding, 1951 |
| <i>A. tritici</i> | Foliar plant parasite | Dried plant material | 9-30 yr | Fielding, 1951 |
| <i>Ditylenchusdipsaci</i> | Foliar plant parasite | Dried plant material | 16- 23 yr | Fielding, 1951 |
| <i>Ditylenchusdipsaci</i> | Foliar plant parasite | -80°C | 5 yr | Cooper <i>et al.</i> , 1971 |
| <i>Filenchuspolyhyhpnus</i> | Foliar in moss | Dried plant material | 39 yr | Steiner and Albin, 1946 |
| <i>Helicotylenchusdihystera</i> | Plant parasite in soil | Dry soil | 250 d | Aroian <i>et al.</i> , 1993 |
| <i>Pratylenchuspenetrans</i> | Plant parasite in soil, roots | Dry soil | 770 d | Townshend, 1984 |

Anoxybiosis: *Bursaphelenchusxylophilus* can survive more than 14 days under anoxic conditions (Kitazume *et al.*, 2018). Many of the soil-inhabiting and intestine-inhabiting nematodes live in habitats that can at times be abnormally low in oxygen content. The nematodes most susceptible to low oxygen levels were those in the process of molting. Physiologically, this is a very active stage in the life cycle of the nematode, during which there is apparently a high demand for oxygen. Oxygen consumption of the surviving infective larvae in the egg was uniform and low. Anaerobic conditions also had pronounced effects on the growth of the developing nematode, egg production, hatching and movement.

References

1. Aroian, RV, Carta L, Alosian, IK and Sternberg, PW, 1993. A free-living *Panagrolaimus* sp. from Armenia can survive in anhydrobiosis for 8.7 years. *J. Nematol*, 25: 500-502.
2. Bird, A F and Bird, J, 1991. *The Structure of Nematodes*. 2nd ed. Academic Press, San Diego, CA pp. 94.
3. Bird, AF and Buttrose, MS, 1974. Ultrastructural changes in the nematode *Anguinatritici* associated with anhydrobiosis. *Journal of Ultrastructure Research*, 48, 177-189.
4. Chapman, RF, 1971. *The Insects. structure and function*. American

- Elsevier Publishing Company, New York.
5. Christie, JR, 1959. Plant Nematodes Their Bionomics and Control. Agricultural Experiment Stations, University of Florida, Gainesville, FL.
 6. Cooper, AFR, and Van Gundy, SD, 1971. Senescence, quiescence, and cryptobiosis. In B. M. Zuckerman, W. F. Mai, and R. A. Rohde [eds.], Plant Parasitic Nematodes. Volume II. Academic Press, New York. pp. 297-318.
 7. Croll, NA, 1970. The behaviour of nematodes: their activity, senses and responses. *The behaviour of nematodes: their activity, senses and responses*. London: Edward Arnold (Publishers) Ltd. pp.117.
 8. Crowe, JH, and Madin, KAC, 1975. Anhydrobiosis in nematodes: evaporative water loss and survival. *Journal of Experimental Zoology*, 193 :323-333.
 9. De Guiran, G, 1979. A necessary diapause in root-knot nematodes. Observations on its distribution and inheritance in *Meloidogyne incognita*. *Revue de Nematology*, 2: 223-231.
 10. Eisenback, JD, 1985. Detailed morphology and anatomy of second-stage juveniles, males, and females of the genus *Meloidogyne* (root-knot nematodes). An advanced treatise on *Meloidogyne*, 1: 47-77.
 11. Ellenby, C, 1946. Nature of the cyst wall of the potato-root eelworm *Heteroderarostochiensis*, Wollenweber, and its permeability to water. *Nature*, 157: 302-303.
 12. Ellenby, C 1968. Desiccation survival in the plant parasitic nematodes, *Heteroderarostochiensis* Wollenweber and *Ditylenchus dipsaci* (Kuhn) Filipjev. *Proceedings of the Royal Society of London. Series B. Biological Sciences*, 169: 203-213.
 13. Ellenby, C, 1969. Dormancy and survival in nematodes. *Symposia of the Society for Experimental Biology*, 23: 572-576.
 14. \Evans, AAF and Perry, RN, 1976. Survival strategies in nematodes, In N.A. Croll [ed.], *The Organization of Nematodes*. Academic Press, London. pp. 383-424.
 15. Fielding, MJ, 1951. Observations on the length of dormancy in certain plant infecting nematodes. *Proceedings of the Helminthological Society of Washington*, 18: 110-112.
 16. Gaur, H.S. & Perry, R.N. (1991). The role of the moulted cuticles in the desiccation survival of adults of *Rotylenchulus reniformis*. *Revue de Nématologie (French)* 14, 491-496.
 17. Gaur, HS, Beane, J and Perry, RN, 2000. The influence of root diffusate, host age and water regimes on hatching of the root-knot nematode, *Meloidogynetriticoryzae*. *Nematology*, 2: 191-199.
 18. Greco, N, 1981. Hatching of *Heteroderacarotae* and *H. avenae*. *Nematologica*, 27: 366-371.
 19. Greco, N, Di Vito, M, and Lamberti, F, 1986. Studies on the biology of *Heteroderagoettingiana* in southern Italy. *Nematologia Mediterranea*, 14: 23-39.
 20. Huang, SP, and Pereira, AC, 1994. Influence of inoculum density, host, and low temperature period on delayed hatch of *Meloidogyne javanica* eggs. *Journal of Nematology*, 26: 72-75.
 21. Ibrahim, SK, Perry, RN, Plowright, RA and Rowe, J, 1993. Hatching behaviour of the rice cyst nematodes *Heteroderasacchari* and *H. oryzicola* in relation to age of host plant. *Fundamental and Applied Nematology*, 16: 23-29.
 22. Ishibashi, N, Kondo, E, Muraoka, M. and Yokoo, T, 1973. Ecological significance of

- dormancy in plant parasitic nematodes I. Ecological difference between eggs in gelatinous matrix and cyst of *Heterodera glycinis* ICHINOHE (Tylenchida: Heteroderidae). *Applied Entomology and Zoology*, 8 :53-63.
23. Kenyon, C, 1997. Environmental factors and gene activities that influence life span. *Cold Spring Harbor Monograph Archive*, pp. 791-813.
 24. Kitazume, H, Dayi, M, Tanaka, R, and Kikuchi, T, 2018. Assessment of the behaviour and survival of nematodes under low oxygen concentrations. *PLoS one*, 13, e0197122.
 25. McClure, MA, Schmitt, ME, and McCullough, MD, 2008. Distribution, biology and pathology of *Anguina pacifica*. *Journal of Nematology*, 40: 226.
 26. Moens, M. and Perry RN, 2009. Migratory plant endoparasitic nematodes: A group rich in contrasts and divergence. *Annual Review of Phytopathology*, 47: 313-332.
 27. Mota, MM, and Vieira, P, 2008. *Pine wilt disease: a worldwide threat to forest ecosystems*. Heidelberg, Germany. Springer, pp. 450.
 28. Needham, JT, 1743. Concerning certain chalky concretions, called malm; with some microscopical observations on the farina of Red Lilly, and worms discovered in Smutty Corn. *Phil Transact Royal Soc Lond*, 42: 634.
 29. Orion, D, and Kritzman, G, 2001. A role of the gelatinous matrix in the resistance of root-knot nematode (*Meloidogyne* spp.) eggs to microorganisms. *Journal of Nematology*, 33: 203.
 30. Perry, RN, and Moens, M, 2011. Survival of parasitic nematodes outside the host. In: Perry, R.N. & Wharton, D.A. (Eds). *Molecular and Physiological Basis of Nematode Survival*. UK, CAB International, pp. 1-27.
 31. Perry, RN 1989. Dormancy and hatching of nematode eggs. *Parasitology Today*, 5: 377-383.
 32. Perry, RN, 2002. Hatching. *The biology of nematodes*, 6: 147-169.
 33. Perry, RN, 2011. Understanding the survival strategies of nematodes. *Animal Science Reviews*, 99-104.
 34. Preston, CM, Bird, AF, 1987. Physiological and morphological changes associated with recovery from anabiosis in the dauer larva of the nematode *Anguina agrotis*. *Parasitology*, 95: 125-133.
 35. Romoser, WS, and Stoffolano, JG, 1998. *The Science of Entomology*, 4th ed. WCB/McGraw-Hill, Boston, MA.
 36. Sikora, EJ, and Noel, GR, 1996. Hatch and emergence of *Heterodera glycinis* in root leachate from resistant and susceptible soybean cultivars. *Journal of Nematology*, 28: 501-509.
 37. Steiner, G, and Albin, FE, 1946. Resuscitation of the nematode *Tylenchus polyhynchus* sp., after almost 39 years' dormancy. *Journal of the Washington Academy of Sciences*, 36: 97-99.
 38. Thompson, JM, and Tylka, GL, 1997. Differences in hatching of *Heterodera glycinis* egg-mass and encysted eggs in vitro. *Journal of Nematology*, 29: 315.
 39. Townshend, JL, 1984. Anhydrobiosis in *Pratylenchus penetrans*. *Journal of Nematology*, 16: 282-289.
 40. Turner, SJ and Rowe, JA, 2006. Cyst nematodes. In: Perry, R. & Moens, M. (Eds). *Plant nematology*. UK, CAB International, pp. 91-122.
 41. Van gundy, SD, 1985. Ecology of *Meloidogyne* spp.- emphasis on

- environmental factors affecting survival and pathogenicity, In J. N. Sasser and C. C. Carter [eds.], *An Advanced Treatise on Meloidogyne*. Volume I. Biology and Control. North Carolina State University Graphics, Raleigh, NC. pp. 177-182.
42. Wesemael, WML, Perry, RN, and Moens, M, 2006. The influence of root diffusate and host age on hatching of the root-knot nematodes, *Meloidogyne chitwoodi* and *M. fallax*. *Nematology*, 8: 895-902.
43. Wharton, DA, 1986. *A Functional Biology of Nematodes*. The Johns Hopkins University Press, Baltimore, MD.
44. Wharton, DA, 1980. Nematode eggshells. *Parasitology*, 81: 447-463.
45. Womersley, CZ, Wharton, DA and Higa, LM, 1998. Survival biology. in R.N. Perry, and D.J. Wright, D.J., eds. *The physiology and biochemistry of free-living and plant-parasitic nematodes*. Wallingford, UK: CAB International. pp. 271-302.
46. Womersley, C, 1978. A comparison of the rate of drying of four nematode species using a liquid paraffin technique. *Annals of Applied Biology*, 90: 401-405.
47. Wright, DJ and Perry, RN, 2006. Reproduction, physiology and biochemistry. In: Perry, R.N. & Moens, M. (Eds). *Plant nematology*. 1st ed. UK, CAB International, pp. 187-209.
48. Yen, JH, Niblack, TL and Wiebold WJ, 1995. Dormancy of *Heterodera glycines* in Missouri. *Journal of Nematology*, 27: 153-163.
49. Zheng, L, and Ferris, H, 1991. Four types of dormancy exhibited by eggs of *Heterodera schachtii*. *Revue de Nematology* 14: 419-426.